

**Unit  
04**

## **ALGEBRAIC EXPRESSIONS AND ALGEBRAIC FORMULAS**

**Define the following terms**

**Algebraic Expressions**

When operations of addition and subtraction are applied to algebraic terms we obtain an algebraic expression. For **Polynomials**.

A polynomial in the variable  $x$  is an algebraic expression of the form

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + a_1 x + a_0, a_n \neq 0, \dots (i)$$

Where  $n$ , the highest power of  $x$ , is a non-negative integer called the degree of the polynomial and each coefficient  $a_n$  is a real number. The coefficient  $a_n$  of the highest power of  $x$  is called the leading coefficient of the polynomial.  $2x^4y^3 + x^2y^2 + 8x$  is a polynomial in two variables  $x$  and  $y$  and has degree 7.

**Rational Express Expression**

The quotient  $\frac{p(x)}{q(x)}$  of two polynomials  $p(x)$  and  $q(x)$ , where  $q(x)$  is a non-zero polynomial, is called a rational expression.

For example,  $\frac{2x+1}{3x+8}$ ,  $3x+8 \neq 0$  is a

rational expression.

In the rational expression  $\frac{p(x)}{q(x)}$ ,  $p(x)$  is called the numerator and  $q(x)$  is known as the denominator of the rational expression  $\frac{p(x)}{q(x)}$ . The rational expression

$\frac{p(x)}{q(x)}$  need not be a polynomial.

instance,  $5x^2 - 3x + \frac{2}{\sqrt{x}}$  and

$3xy + \frac{3}{x}$  ( $x \neq 0$ ) are algebraic expressions.

**Example**

Reduce the following algebraic fractions to their lowest forms.

$$(i) \quad \frac{lx+mx-ly-my}{3x^2-3y^2} \quad (ii) \quad \frac{3x^2+18x+27}{5x^2-45}$$

**Solution**

$$\begin{aligned} (i) \quad & \frac{lx+mx-ly-my}{3x^2-3y^2} \\ &= \frac{x(l+m)-y(l+m)}{3(x^2-y^2)} \\ &= \frac{(l+m)(x-y)}{3(x+y)(x-y)} \\ &= \frac{l+m}{3(x+y)} \end{aligned}$$

Which is in the lowest forms.

$$(ii) \frac{3x^2 + 18x + 27}{5x^2 - 45} = \frac{3(x^2 + 6x + 9)}{5(x^2 - 9)}$$
$$\frac{3(x+3)(x+3)}{5(x+3)(x-3)}$$

$$\frac{3(x+3)}{5(x-3)}$$

Which is in the lowest forms

### Example

Simplify (i)  $\frac{1}{x-y} - \frac{1}{x+y} + \frac{2x}{x^2-y^2}$

(ii)  $\frac{2x^2}{x^4-16} - \frac{x}{x^2-4} + \frac{1}{x+2}$

### Solution

$$(i) \frac{1}{x-y} - \frac{1}{x+y} + \frac{2x}{x^2-y^2}$$
$$= \frac{1}{x-y} - \frac{1}{x+y} + \frac{2x}{(x+y)(x-y)}$$
$$= \frac{x+y-(x-y)+2x}{(x+y)(x-y)}$$

(L.C.M of denominators)

$$= \frac{x+y-x+y+2x}{(x+y)(x-y)}$$
$$= \frac{2x+2y}{(x+y)(x-y)}$$

$$= \frac{2(x+y)}{(x+y)(x-y)} = \frac{2}{x-y}$$

$$(ii) \frac{2x^2}{x^4-16} - \frac{x}{x^2-4} + \frac{1}{x+2}$$

$$= \frac{2x^2}{(x^2+4)(x^2-4)} - \frac{x}{x^2-4} + \frac{1}{x+2}$$

$$= \frac{2x^2}{(x^2+4)(x+2)(x-2)} - \frac{x}{(x+2)(x-2)} + \frac{1}{x+2}$$

$$= \frac{2x^2 - x(x^2+4) + (x^2+4)(x-2)}{(x^2+4)(x+2)(x-2)} = \frac{2x^2 - x^3 - 4x^2 - x^3 + 4x^2 - 8}{(x^2+4)(x+2)(x-2)}$$

$$= \frac{-8}{(x^2+4)(x+2)(x-2)}$$

$$= \frac{-8}{(x^2+4)(x^2-4)} = \frac{-8}{x^4-16}$$

**Example**

Find the product  $\frac{x+2}{2x-3y} \cdot \frac{4x^2-9y^2}{xy+2y}$

**Solution**

$$\begin{aligned} \frac{x+2}{2x-3y} \cdot \frac{4x^2-9y^2}{xy+2y} &= \frac{(x+2)[(2x)^2-(3y)^2]}{(2x-3y)(x+2)y} \\ &= \frac{(x+2)(2x+3y)(2x-3y)}{y(x+2)(2x-3y)} \\ &= \frac{2x+3y}{y} \end{aligned}$$

**Example**

Simplify  $\frac{7xy}{x^2-4x+4} \div \frac{14y}{x^2-4}$

**Solution**

$$\begin{aligned} &\frac{7xy}{x^2-4x+4} \div \frac{14y}{x^2-4} \\ &= \frac{7xy}{x^2-4x+4} \times \frac{x^2-4}{14y} \end{aligned}$$

$$\begin{aligned} &= \frac{7xy}{(x-2)(x+2)} \times \frac{(x+2)(x-2)}{14y} \\ &= \frac{x(x+2)}{2(x-2)} \end{aligned}$$

**Example**

Evaluate  $\frac{3x^2\sqrt{y+6}}{5(x+y)}$  if  $x = -4$  and  $y = 9$

**Solution**

We have, by putting  $x = -4$  and  $y = 9$ ,

$$\frac{3x^2\sqrt{y+6}}{5(x+y)} = \frac{3(-4)^2\sqrt{9+6}}{5(-4+9)} = \frac{3(16)\sqrt{15}}{5(5)} = \frac{150}{25} = 6$$

## Exercise 4.1

1. Identify whether the following algebraic expression are polynomials (yes or no).

- (i)  $3x^2 + \frac{1}{x} - 5$  No
- (ii)  $3x^3 - 4x^2 - x\sqrt{x} + 3$  No
- (iii)  $x^2 - 3x + \sqrt{2}$  Yes
- (iv)  $\frac{3x}{2x-1} + 8$  No

2. State whether each of the following expression is a rational expression or not.

- (i)  $\frac{3\sqrt{x}}{3\sqrt{x}+5}$  No
- (ii)  $\frac{x^3 - 2x^2 + \sqrt{3}}{2+3x-x^2}$  Yes
- (iii)  $\frac{x^2 + 6x + 9}{x^2 - 9}$  Yes

$$(iv) \quad \frac{2\sqrt{x}+3}{2\sqrt{x}-3} \quad \text{No}$$

3. Reduce the following rational expression to the lowest forms.

$$(i) \quad \frac{120x^2y^3z^5}{30x^3yz^2}$$

$$= 4x^{2-3}y^{3-1}z^{5-2}$$

$$= 4x^{-1}y^2z^3$$

$$= \frac{4y^2z^3}{x}$$

$$(ii) \quad \frac{8a(x+1)}{2(x^2-1)} = \frac{4a(x+1)}{(x-1)(x+1)} = \frac{4a}{x-1}$$

$$(iii) \quad \frac{(x+y)^2 - 4xy}{(x-y)^2} = \frac{x^2 + y^2 + 2xy - 4xy}{(x-y)(x-y)}$$

$$= \frac{x^2 + y^2 - 2xy}{(x-y)(x-y)}$$

$$= \frac{(x-y)^2}{(x-y)(x-y)}$$

$$= \frac{\cancel{(x-y)^2}}{\cancel{(x-y)^2}} = 1$$

$$(iv) \quad \frac{(x^3 - y^3)(x^2 - 2xy + y^2)}{(x-y)(x^2 + xy + y^2)}$$

$$= \frac{\cancel{(x^3 - y^3)}(x-y)^2}{\cancel{x^3 - y^3}} = (x-y)^2$$

$$(v) \quad \frac{(x+2)(x^2-1)}{(x+1)(x^2-4)}$$

$$= \frac{\cancel{(x+2)}(x-1)\cancel{(x+1)}}{\cancel{(x+1)}(x-2)\cancel{(x+2)}} = \frac{x-1}{x-2}$$

$$(vi) \quad \frac{x^2 - 4x + 4}{2x^2 - 8} = \frac{(x-2)^2}{2(x^2-4)}$$

$$= \frac{(x-2)^2}{2(x-2)(x+2)}$$

$$= \frac{\cancel{(x-2)}(x-2)}{\cancel{2(x-2)}(x+2)}$$

$$= \frac{x-2}{2(x+2)}$$

$$(vii) \quad \frac{64x^5 - 64x}{(8x^2 + 8)(2x+2)}$$

$$= \frac{64x(x^4 - 1)}{8(x^2 + 1).2(x+1)}$$

$$= \frac{64x(x^4 - 1)}{16(x^2 + 1)(x+1)}$$

$$= \frac{4x(x^2 + 1)(x^2 - 1)}{(x^2 + 1)(x+1)}$$

$$= \frac{4x \cancel{(x^2 + 1)} (x-1) \cancel{(x+1)}}{\cancel{(x^2 + 1)} \cancel{(x+1)}}$$

$$= 4x(x-1)$$

$$\frac{9x^2 - (x^2 - 4)^2}{4 + 3x - x^2} = \frac{(3x)^2 - (x^2 - 4)^2}{4 + 3x - x^2}$$

$$= \frac{(3x + x^2 - 4) \cancel{(3x - x^2 + 4)}}{\cancel{(4 + 3x - x^2)}}$$

$$= 3x + x^2 - 4$$

$$= x^2 + 3x - 4$$

4. Evaluate (a)  $\frac{x^3y - 2z}{xz}$  for (i)  $x = 3$

$$y = -1, z = -2.$$

$$(a) \quad \frac{(3)^3(-1) - 2(-2)}{3(-2)} = \frac{-27 + 4}{-6}$$

$$= \frac{-23}{-6} = \frac{23}{6} = 3\frac{5}{6}$$

$$(b) \frac{x^2y^3 - 5z^4}{xyz} \text{ for } x = 4, y = -2, z =$$

-1

$$= \frac{(4)^2(-2)^3 - 5(-1)^4}{(4)(-2)(-1)} = \frac{-16(8) - 5}{8}$$

$$= \frac{-128 - 5}{8} = \frac{-133}{8} = -16\frac{5}{8}$$

5. Perform the indicated operation and simplify

$$(i) \frac{15}{2x-3y} - \frac{4}{3y-2x}$$

$$= \frac{15(3y-2x) - 4(2x-3y)}{(2x-3y)(3y-2x)}$$

$$= \frac{45y - 30x - 8x + 12y}{(2x-3y)(3y-2x)}$$

$$= \frac{57y - 38x}{(2x-3y)(3y-2x)}$$

$$= \frac{19(3y-2x)}{(2x-3y)(3y-2x)} = \frac{19}{2x-3y}$$

$$(ii) \frac{1+2x}{1-2x} - \frac{1-2x}{1+2x}$$

$$= \frac{(1+2x)^2 - (1-2x)^2}{(1-2x)(1+2x)}$$

$$= \frac{(1+4x^2 + 4x) - (1+4x^2 - 4x)}{(1-2x)(1+2x)}$$

$$= \frac{1+4x^2 + 4x - 1-4x^2 + 4x}{(1-2x)(1+2x)}$$

$$= \frac{8x}{(1-2x)(1+2x)} = \frac{8x}{1-4x^2}$$

$$(iii) \frac{x^2 - 25}{x^2 - 36} - \frac{x+5}{x+6}$$

$$= \frac{(x-5)(x+5)}{(x-6)(x+6)} - \frac{x+5}{x+6}$$

$$= \frac{(x-5)(x+5) - (x+5)(x-6)}{(x+6)(x-6)}$$

$$= \frac{(x+5)[(x-5) - (x-6)]}{(x+6)(x-6)}$$

$$= \frac{(x+5)(\cancel{x-5} - \cancel{x+6})}{(x+6)(x-6)}$$

$$= \frac{(x+5)(1)}{(x+6)(x-6)} = \frac{x+5}{x^2 - 36}$$

$$(iv) \frac{x}{x-y} - \frac{y}{x+y} - \frac{2xy}{x^2 - y^2}$$

$$= \frac{x(x+y) - y(x-y)}{(x-y)(x+y)} - \frac{2xy}{x^2 - y^2}$$

$$= \frac{x^2 + xy - \cancel{xy} + y^2}{x^2 - y^2} - \frac{2xy}{x^2 - y^2}$$

$$= \frac{x^2 + y^2 - 2xy}{x^2 - y^2}$$

$$= \frac{x^2 + y^2 - 2xy}{(x^2 - y^2)}$$

$$= \frac{(x-y)^2}{(\cancel{x-y})(x+y)} = \frac{x-y}{x+y}$$

$$(v) \frac{x-2}{x^2 + 6x + 9} - \frac{x+2}{2x^2 - 18}$$

$$= \frac{x-2}{x^2 + 3x + 3x + 9} - \frac{x+2}{2(x^2 - 9)}$$

$$= \frac{x-2}{x(x+3) + 3(x+3)} - \frac{x+2}{2(x-3)(x+3)}$$

$$= \frac{x-2}{(x+3)(x+3)} - \frac{x+2}{2(x-3)(x+3)}$$

$$= \frac{2(x-3)(x-2) - (x+3)(x+2)}{2(x-3)(x+3)(x+3)}$$

$$= \frac{2(x^2 - 2x - 3x + 6) - (x^2 + 2x + 3x + 6)}{2(x-3)(x+3)^2}$$

$$\begin{aligned}
 &= \frac{2(x^2 - 5x + 6) - (x^2 + 5x + 6)}{2(x-3)(x+3)^2} \\
 &= \frac{2x^2 - 10x + 12 - x^2 - 5x - 6}{2(x-3)(x+3)^2} \\
 &= \frac{x^2 - 15x + 6}{2(x-3)(x+3)^2}
 \end{aligned}$$

$$\begin{aligned}
 (\text{vi}) \quad & \frac{1}{x-1} - \frac{1}{x+1} - \frac{2}{x^2+1} - \frac{4}{x^4-1} \\
 &= \frac{x+1-(x-1)}{(x-1)(x+1)} - \frac{2}{x^2+1} - \frac{4}{x^4-1} \\
 &= \frac{x+1-x+1}{x^2-1} - \frac{2}{x^2+1} - \frac{4}{x^4-1} \\
 &= \frac{2}{x^2-1} - \frac{2}{x^2+1} - \frac{4}{x^4-1} \\
 &= \frac{2(x^2+1)-2(x^2-1)}{(x^2-1)(x^2+1)} - \frac{4}{x^4-1} \\
 &= \frac{2x^2+2-2x^2+2}{x^4-1} - \frac{4}{x^4-1} \\
 &= \frac{4}{x^4-1} - \frac{4}{x^4-1} \\
 &= \frac{4-4}{x^4-1} \\
 &= \frac{0}{x^4-1} \\
 &= 0
 \end{aligned}$$

**6. Perform the indicated operation and simplify:**

$$\begin{aligned}
 (\text{i}) \quad & (x^2 - 49) \frac{5x+2}{x+7} \\
 &= (x-7)(\cancel{x+7}) \frac{5x+2}{\cancel{x+7}} \\
 &= (x-7)(5x+2)
 \end{aligned}$$

$$\begin{aligned}
 (\text{ii}) \quad & \frac{4x-12}{x^2-9} \div \frac{18-2x^2}{x^2+6x+9} \\
 &= \frac{4(x-3)}{(x-3)(x+3)} \div \frac{2(9-x^2)}{x^2+3x+3x+9} \\
 &= \frac{4(x-3)}{(x-3)(x+3)} \div \frac{2(3-x)(3+x)}{x(x+3)+3(x+3)} \\
 &= \frac{4(x-3)}{(x-3)(x+3)} \div \frac{2(3-x)(3+x)}{(x+3)(x+3)} \\
 &= \frac{4(x-3)}{(x+3)(x-3)} \times \frac{(x+3)(x+3)}{2(3+x)(3-x)} \\
 &= \frac{2}{3-x} \\
 (\text{iii}) \quad & \frac{x^6-y^6}{x^2-y^2} \div (x^4+x^2y^2+y^4) \\
 &= \frac{(x^3)^2-(y^3)^2}{x^2-y^2} \div (x^4+x^2y^2+y^4) \\
 &= \frac{(x^3-y^3)(x^3+y^3)}{x^2-y^2} \div (x^4+x^2y^2+y^4) \\
 &= \frac{(x-y)(x^2+xy+y^2)(x+y)(x^2-xy+y^2)}{x^2-y^2} \\
 &\times \frac{1}{x^4+x^2y^2+y^4} \\
 &= \frac{\cancel{(x^2-y^2)}(x^2+xy+y^2)(x^2-xy+y^2)}{\cancel{x^2-y^2}} \\
 &\times \frac{1}{x^4+x^2y^2+y^4} \\
 &= \frac{x^4+x^2y^2+y^4}{x^4+x^2y^2+y^4} = 1 \\
 (\text{iv}) \quad & \frac{x^2-1}{x^2+2x+1} \cdot \frac{x+5}{1-x} \\
 &= \frac{-(x-1)(x+1)}{x^2+x+x+1} \cdot \frac{x+5}{(x-1)}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{-(x+1)(x+5)}{x(x+1)+1(x+1)} \\
 &= \frac{-(x+1)(x+5)}{(x+1)(x+1)} = -\frac{x+5}{x+1} \\
 (\text{v}) \quad &\frac{x^2+xy}{y(x+y)} \cdot \frac{x^2+xy}{y(x+y)} + \frac{x^2-x}{xy-2y} \\
 &= \frac{x(x+y)}{y(x+y)} \cdot \frac{x(x+y)}{y(x+y)} \times \frac{y(x-2)}{x(x-1)} \\
 &= \frac{x(x-2)}{y(x-1)}
 \end{aligned}$$

**Example**

If  $a+b=7$  and  $a-b=3$ , then find the value of (a)  $a^2+b^2$  (b)  $ab$

**Solution**

We are given that  $a+b=7$  and  $a-b=3$

(a) To find the value of  $(a^2+b^2)$ , we use the formula

$$(a+b)^2 + (a-b)^2 = 2(a^2+b^2)$$

Substituting the values  $a+b=7$  and  $a-b=3$ , we get

$$\begin{aligned}
 (7)^2 + (3)^2 &= 2(a^2+b^2) \\
 \Rightarrow 49+9 &= 2(a^2+b^2) \\
 \Rightarrow 58 &= 2(a^2+b^2), \\
 \Rightarrow 29 &= a^2+b^2,
 \end{aligned}$$

(b) To find the value of  $ab$ , we make use of the formula

$$\begin{aligned}
 (a+b)^2 - (a-b)^2 &= 4ab \\
 \Rightarrow (7)^2 - (3)^2 &= 4ab,
 \end{aligned}$$

$$\begin{aligned}
 \Rightarrow 49-9 &= 4ab \\
 \Rightarrow 40 &= 4ab, \\
 \Rightarrow 10 &= ab,
 \end{aligned}$$

Hence  $a^2+b^2=29$  and  $ab=10$ .

**Example**

If  $a^2+b^2+c^2=43$  and  $ab+bc+ca=3$ , then find the value of  $a+b+c$ .

**Solution**

We know that

$$(a+b+c)^2 = a^2+b^2+c^2+2ab+2bc+2ca$$

$$(a+b+c)^2 = a^2+b^2+c^2+2(ab+bc+ca)$$

$$\Rightarrow (a+b+c)^2 = 43+2 \times 3$$

(Putting  $a^2+b^2+c^2=43$  and  $ab+bc+ca=3$ )

$$\Rightarrow (a+b+c)^2 = 49$$

$$\Rightarrow a+b+c = \pm \sqrt{49}$$

Hence  $a+b+c = \pm 7$

**Example**

If  $a+b+c=6$  and  $a^2+b^2+c^2=24$  then find the value of  $ab+bc+ca$ .

**Solution**

We have

$$(a+b+c)^2 = a^2+b^2+c^2+2ab+2bc+2ca$$

$$(6)^2 = 24+2(ab+bc+ca)$$

$$\Rightarrow 36 = 24+2(ab+bc+ca)$$

$$\Rightarrow 12 = 2(ab+bc+ca)$$

Hence  $ab+bc+ca=6$

**Example**

If  $a+b+c=7$  and  $ab+bc+ca=9$ , then find the value of  $a^2+b^2+c^2$ .

**Solution**

We know that

$$\begin{aligned}(a+b+c)^2 &= a^2 + b^2 + c^2 + 2ab + 2bc + 2ca \\ \Rightarrow (a+b+c)^2 &= a^2 + b^2 + c^2 + 2(ab+bc+ca) \\ \Rightarrow (7)^2 &= a^2 + b^2 + c^2 + 2(9) \\ \Rightarrow 49 &= a^2 + b^2 + c^2 + 18 \\ \Rightarrow 31 &= a^2 + b^2 + c^2 \\ \text{Hence } a^2 + b^2 + c^2 &= 31\end{aligned}$$

**Example**

If  $2x - 3y = 10$  and  $xy = 2$ , then find the value of  $8x^3 - 27y^3$ .

**Solution**

We are given that  $2x - 3y = 10$

$$\begin{aligned}\Rightarrow (2x-3y)^3 &= (10)^3 \\ \Rightarrow 8x^3 - 27y^3 - 3 \times 2x \times 3y(2x-3y) &= 1000 \\ \Rightarrow 8x^3 - 27y^3 - 18xy(2x-3y) &= 1000 \\ \Rightarrow 8x^3 - 27y^3 - 18 \times 2 \times 10 &= 1000 \\ \Rightarrow 8x^3 - 27y^3 - 360 &= 1000 \\ \text{Hence } 8x^3 - 27y^3 &= 1360\end{aligned}$$

**Example**

If  $x + \frac{1}{x} = 8$ , then find the value of  $x^3 + \frac{1}{x^3}$

**Solution**

We have been given  $x + \frac{1}{x} = 8$

$$\begin{aligned}\Rightarrow \left(x + \frac{1}{x}\right)^3 &= (8)^3 \\ \Rightarrow x^3 + \frac{1}{x^3} + 3 \times x \times \frac{1}{x} \left(x + \frac{1}{x}\right) &= 512\end{aligned}$$

$$\begin{aligned}\Rightarrow x^3 + \frac{1}{x^3} + 3 \left(x + \frac{1}{x}\right) &= . \quad 512 \\ \Rightarrow x^3 + \frac{1}{x^3} + 3 \times 8 &= \quad 512 \\ \Rightarrow x^3 + \frac{1}{x^3} + 24 &= \quad 512 \\ \Rightarrow x^3 + \frac{1}{x^3} &= 512 - 24 \\ \Rightarrow x^3 + \frac{1}{x^3} &= 488\end{aligned}$$

**Example**

If  $x - \frac{1}{x} = 4$ , then find  $x^3 - \frac{1}{x^3}$

**Solution**

We have  $x - \frac{1}{x} = 4$

$$\begin{aligned}\Rightarrow \left(x - \frac{1}{x}\right)^3 &= (4)^3 \\ \Rightarrow x^3 - \frac{1}{x^3} - 3x \times \frac{1}{x} \left(x - \frac{1}{x}\right) &= 64 \\ \Rightarrow x^3 - \frac{1}{x^3} - 3(4) &= 64 \\ \Rightarrow x^3 - \frac{1}{x^3} - 12 &= 64 \\ \Rightarrow x^3 - \frac{1}{x^3} &= 64 + 12 \\ \Rightarrow x^3 - \frac{1}{x^3} &= 76\end{aligned}$$

**Example**

Factorize  $64x^3 + 343y^3$

**Solution**

We have

$$64x^3 + 343y^3 = (4x)^3 + (7y)^3$$

$$= (4x+7y)[(4x)^2 - (4x)(7y) + (7y)^2]$$

$$= (4x+7y)(16x^2 - 28xy + 49y^2)$$

**Example**

Factorize  $125x^3 - 1331y^3$

**Solution**

We have

$$125x^3 - 1331y^3 = (5x)^3 - (11y)^3$$

$$= (5x - 11y)[(5x)^2 + (5x)(11y) + (11y)^2]$$

$$= (5x - 11y)(25x^2 + 55xy + 121y^2)$$

**Example**

Factorize

$$\left(\frac{2}{3}x + \frac{3}{2x}\right)\left(\frac{4}{9}x^2 - 1 + \frac{9}{4x^2}\right)$$

**Solution**

$$\left(\frac{2}{3}x + \frac{3}{2x}\right)\left(\frac{4}{9}x^2 - 1 + \frac{9}{4x^2}\right)$$

$$= \left(\frac{2}{3}x + \frac{3}{2x}\right)\left[\left(\frac{2}{3}x\right)^2 - \left(\frac{2}{3}x\right)\left(\frac{3}{2x}\right) + \left(\frac{3}{2x}\right)^2\right]$$

$$= \left(\frac{2}{3}x\right)^3 + \left(\frac{3}{2x}\right)^3$$

$$= \frac{8}{27}x^3 + \frac{27}{8x^3}$$

**Example**

Find the product  $\left(\frac{4}{5}x - \frac{5}{4x}\right)\left(\frac{16}{25}x^2 + \frac{25}{16x^2} + 1\right)$

**Solution**

$$\left(\frac{4}{5}x - \frac{5}{4x}\right)\left(\frac{16}{25}x^2 + \frac{25}{16x^2} + 1\right)$$

$$= \left(\frac{4}{5}x - \frac{5}{4x}\right)\left(\frac{16x^2}{25} + 1 + \frac{25}{16x^2}\right)$$

(rearranging)

$$= \left(\frac{4}{5}x - \frac{5}{4x}\right)\left[\left(\frac{4}{5}x\right)^2 + \left(\frac{4}{5}x\right)\left(\frac{5}{4x}\right) + \left(\frac{5}{4x}\right)^2\right]$$

$$= \left(\frac{4}{5}x\right)^3 - \left(\frac{5}{4x}\right)^3 = \frac{64}{125}x^3 - \frac{125}{64x^3}$$

**Example**

Find the continued product of  $(x+y)(x-y)(x^2+xy+y^2)(x^2-xy+y^2)$

**Solution**

$$(x+y)(x-y)(x^2+xy+y^2)(x^2-xy+y^2)$$

$$= (x+y)(x^2-xy+y^2)(x-y)(x^2+xy+y^2)$$

$$= (x^3+y^3)(x^3-y^3) = (x^3)^2 - (y^3)^2 = x^6 - y^6$$

## Exercise 4.2

1.(i) If  $a+b=10$  and  $a-b=6$  then find value of  $a^2+b^2$ .

**Solution:**

$$2(a^2+b^2) = (a+b)^2 + (a-b)^2$$

$$2(a^2+b^2) = (10)^2 + (6)^2$$

$$2(a^2+b^2) = 100 + 36$$

$$a^2 + b^2 = \frac{136}{2} = 68$$

(ii) If  $a+b=5$ ,  $a-b=\sqrt{17}$  then find value of  $ab$ .

**Solution:**

$$4ab = (a+b)^2 - (a-b)^2$$

$$4ab = (5)^2 - (\sqrt{17})^2$$

$$4ab = 25 - 17$$

$$4ab = 8$$

$$ab = \frac{8}{4} = 2$$

2. If  $a^2 + b^2 + c^2 = 45$  and  $a + b + c = -1$  find value of  $ab + bc + ca$ .

**Solution:**

$$a+b+c = -1$$

Squaring

$$(a+b+c)^2 = (-1)^2$$

$$a^2 + b^2 + c^2 + 2ab + 2bc + 2ca = 1$$

$$a^2 + b^2 + c^2 + 2(ab + bc + ca) = 1$$

$$45 + 2(ab + bc + ca) = 1$$

$$2(ab + bc + ca) = 1 - 45$$

$$2(ab + bc + ca) = -44$$

$$ab + bc + ca = \frac{-44}{2} = -22$$

3. If  $m+n+p = 10$ ,  $mn + np + pm = 27$  find value of  $m^2 + n^2 + p^2$ .

**Solution:**

$$m+n+p = 10$$

Squaring both sides

$$(m+n+p)^2 = (10)^2$$

$$m^2 + n^2 + p^2 + 2mn + 2np + 2mp = 100$$

$$m^2 + n^2 + p^2 + 2(mn + np + mp) = 100$$

$$m^2 + n^2 + p^2 + 2(27) = 100$$

$$m^2 + n^2 + p^2 + 54 = 100$$

$$m^2 + n^2 + p^2 = 100 - 54$$

$$m^2 + n^2 + p^2 = 46$$

4. If  $x^2 + y^2 + z^2 = 78$  and  $y+yz+zx=59$  find  $x + y + z$ .

**Solution:**

$$\begin{aligned}(x+y+z)^2 &= x^2 + y^2 + z^2 + 2xy + 2yz + 2zx \\&= x^2 + y^2 + z^2 + 2(xy + yz + zx) \\&= 78 + 2(59) \\&= 78 + 118 \\&= 196\end{aligned}$$

$$\sqrt{(x+y+z)^2} = \sqrt{196} = \sqrt{(\pm 14)^2}$$

$$x + y + z = \pm 14$$

5. If  $x + y + z = 12$  and  $x^2 + y^2 + z^2 = 64$  find value of  $xy + yz + zx$ .

**Solution:**

$$x + y + z = 12$$

Squaring both sides

$$(x + y + z)^2 = (12)^2$$

$$x^2 + y^2 + z^2 + 2xy + 2yz + 2zx = 144$$

$$x^2 + y^2 + z^2 + 2(xy + yz + zx) = 144$$

$$64 + 2(xy + yz + zx) = 144$$

$$2(xy + yz + zx) = 144 - 64$$

$$2(xy + yz + zx) = 80$$

$$xy + yz + zx = \frac{80}{2} = 40.$$

6. If  $x + y = 7$  and  $xy = 12$  then find value of  $x^3 + y^3$ .

**Solution:**

$$x + y = 7$$

$$(x + y)^3 = (7)^3$$

$$x^3 + y^3 + 3xy(x+y) = 343$$

$$x^3 + y^3 + 3(12)(7) = 343$$

$$x^3 + y^3 + 252 = 343$$

$$x^3 + y^3 = 343 - 252$$

$$x^3 + y^3 = 91$$

7. If  $3x + 4y = 11$  and  $xy = 12$  then find value of  $27x^3 + 64y^3$ .

**Solution:**

$$3x + 4y = 11$$

$$(3x + 4y)^3 = (11)^3$$

$$(3x)^3 + (4y)^3 + 3(3x)(4y)(3x+4y) =$$

$$1331$$

$$27x^3 + 64y^3 + 36xy(3x+4y) = 1331$$

$$27x^3 + 64y^3 + 36(12)(11) = 1331$$

$$27x^3 + 64y^3 + 4752 = 1331$$

$$27x^3 + 64y^3 = 1331 - 4752 = -3421$$

8. If  $x - y = 4$  and  $xy = 21$  then find value of  $x^3 - y^3$ .

**Solution:**

$$x - y = 4$$

$$(x-y)^3 = (4)^3$$

$$x^3 - y^3 - 3xy(x-y) = 64$$

$$x^3 - y^3 - 3(21)(4) = 64$$

$$x^3 - y^3 - 252 = 64$$

$$x^3 - y^3 = 64 + 252$$

$$x^3 - y^3 = 316$$

9. If  $5x - 6y = 13$  and  $xy = 6$  then find value of  $125x^3 - 216y^3$ .

Solution:

$$5x - 6y = 13$$

$$\Rightarrow (5x-6y)^3 = (13)^3$$

$$\Rightarrow (5x)^3 - (6y)^3 - 3(5x)(6y)(5x-6y) = 2197$$

$$125x^3 - 216y^3 - 90xy(5x-6y) = 2197$$

$$125x^3 - 216y^3 - 90(6)(13) = 2197$$

$$125x^3 - 216y^3 - 7020 = 2197$$

$$125x^3 - 216y^3 = 2197 + 7020$$

$$125x^3 - 216y^3 = 9217$$

10. If  $x + \frac{1}{x} = 3$  then find  $x^3 + \frac{1}{x^3}$ .

$$x + \frac{1}{x} = 3 \text{ Cubing both sides}$$

$$\left(x + \frac{1}{x}\right)^3 = (3)^3$$

$$x^3 + \frac{1}{x^3} + 3\left(x\right)\left(\frac{1}{x}\right)\left(x + \frac{1}{x}\right) = 27$$

$$x^3 + \frac{1}{x^3} + 3\left(x + \frac{1}{x}\right) = 27$$

$$x^3 + \frac{1}{x^3} + 3(3) = 27$$

$$x^3 + \frac{1}{x^3} = 27 - 9$$

$$x^3 + \frac{1}{x^3} = 18$$

11. If  $x - \frac{1}{x} = 7$ , then find value of

$$x^3 - \frac{1}{x^3}$$

$$x - \frac{1}{x} = 7 \text{ Taking cube of both sides}$$

$$\left(x - \frac{1}{x}\right)^3 = (7)^3$$

$$x^3 - \frac{1}{x^3} - 3\left(x\right)\left(\frac{1}{x}\right)\left(x - \frac{1}{x}\right) = 343$$

$$x^3 - \frac{1}{x^3} - 3\left(x - \frac{1}{x}\right) = 343$$

$$x^3 - \frac{1}{x^3} - 3(7) = 343$$

$$x^3 - \frac{1}{x^3} - 21 = 343$$

$$x^3 - \frac{1}{x^3} = 343 + 21$$

$$x^3 - \frac{1}{x^3} = 364$$

12. If  $3x + \frac{1}{3x} = 5$ , then find value of

$$27x^3 + \frac{1}{27x^3}$$

$$\left(3x + \frac{1}{3x}\right)^3 = (5)^3$$

$$(3x)^3 + \left(\frac{1}{3x}\right)^3 + 3(3x)\left(\frac{1}{3x}\right)\left(3x + \frac{1}{3x}\right) = 125$$

$$27x^3 + \frac{1}{27x^3} + 3\left(3x + \frac{1}{3x}\right) = 125$$

$$27x^3 + \frac{1}{27x^3} + 3(5) = 125$$

$$27x^3 + \frac{1}{27x^3} + 15 = 125$$

$$27x^3 + \frac{1}{27x^3} = 125 - 15$$

$$27x^3 + \frac{1}{27x^3} = 110$$

13. If  $\left(5x - \frac{1}{5x}\right) = 6$ , then find value of

$$125x^3 - \frac{1}{25x^3}.$$

$$\left(5x - \frac{1}{5x}\right) = 6$$

Taking cube of both sides

$$\left(5x - \frac{1}{5x}\right)^3 = (6)^3$$

$$(5x)^3 - \left(\frac{1}{5x}\right)^3 - 3\left(5x\right)\left(\frac{1}{5x}\right)\left(5x - \frac{1}{5x}\right) = 216$$

$$125x^3 - \frac{1}{125x^3} - 3\left(5x - \frac{1}{5x}\right) = 216$$

$$125x^3 - \frac{1}{125x^3} - 3(6) = 216$$

$$125x^3 - \frac{1}{25x^3} - 18 = 216$$

$$125x^3 - \frac{1}{125x^3} = 216 + 18$$

$$125x^3 - \frac{1}{125x^3} = 234$$

14. Factorize (i)  $x^3 - y^3 - x + y$

$$(i) \quad x^3 - y^3 - x + y$$

$$= (x - y)(x^2 + xy + y^2) - 1(x - y)$$

$$= (x - y)[x^2 + xy + y^2 - 1]$$

$$(ii) \quad 8x^3 - \frac{1}{27y^3}$$

$$= (2x)^3 - \left(\frac{1}{3y}\right)^3$$

$$= \left(2x - \frac{1}{3y}\right) \left( (2x)^2 + (2x)\left(\frac{1}{3y}\right) + \left(\frac{1}{3y}\right)^2 \right)$$

$$= \left(2x - \frac{1}{3y}\right) \left( 4x^2 + \frac{2x}{3y} + \frac{1}{9y^2} \right)$$

15. Find products, using formulae

$$(i) \quad (x^2 + y^2)(x^4 - x^2y^2 + y^4)$$

$$= (x^2)^3 + (y^2)^3$$

$$\text{Ref. } (a + b)(a^2 - ab + b^2) = a^3 + b^3$$

$$= x^6 + y^6$$

$$(ii) \quad (x^3 - y^3)(x^6 + x^3y^3 + y^6)$$

$$= (x^3)^3 - (y^3)^3$$

$$\text{Ref. } (a - b)(a^2 + ab + b^2) = a^3 - b^3$$

$$= x^9 - y^9$$

$$(iii) \quad (x - y)(x + y)(x^2 + y^2)(x^2 + xy + y^2)$$

$$(x^2 - xy + y^2)(x^4 - x^2y^2 + y^4)$$

$$= (x - y)(x^2 + xy + y^2)(x + y)(x^2 - xy + y^2)$$

$$(x^2 + y^2)(x^4 - x^2y^2 + y^4)$$

$$= (x^3 - y^3)(x^3 + y^3)[(x^2)^3 + (y^2)^3]$$

$$= [(x^3)^2 - (y^3)^2](x^6 + y^6)$$

$$= (x^6 - y^6)(x^6 + y^6)$$

$$= (x^6)^2 - (y^6)^2$$

$$= x^{12} - y^{12}$$

$$16. \quad (2x^2 - 1)(2x^2 + 1)(4x^4 + 2x^2 + 1)$$

$$(4x^4 - 2x^2 + 1)$$

$$= (2x^2 - 1)(4x^4 + 2x^2 + 1)(2x^2 + 1)$$

$$(4x^4 - 2x^2 + 1)$$

$$= ((2x^2)^3 - (1)^3)((2x^2)^3 + (1)^3)$$

$$\begin{aligned}
 &= (8x^6 - 1)(8x^6 + 1) \\
 &= (8x^6)^2 - (1)^2
 \end{aligned}
 \quad \quad \quad
 \begin{aligned}
 &= 64x^{12} - 1
 \end{aligned}$$

### Define Surd

An irrational radical with rational radicand is called a surd.

Hence the radical  $\sqrt[n]{a}$  is a surd if

- (i)  $a$  is rational
- (ii) the result  $\sqrt[n]{a}$  is irrational.

e.g.,  $\sqrt{3}, \sqrt{2/5}, \sqrt[3]{7}, \sqrt[4]{10}$  are surds.

But  $\sqrt{\pi}$  is not surd because  $\pi$  is not rational.

**Note:** Every surd is an irrational number but every irrational number is not surd

### Example

Simplify by combining similar terms.

$$(i) \quad 4\sqrt{3} - 3\sqrt{27} + 2\sqrt{75}$$

$$(ii) \quad \sqrt[3]{128} - \sqrt[3]{250} + \sqrt[3]{432}$$

### Solution

$$\begin{aligned}
 (i) \quad &4\sqrt{3} - 3\sqrt{27} + 2\sqrt{75} \\
 &= 4\sqrt{3} - 3\sqrt{9 \times 3} + 2\sqrt{25 \times 3} = 4\sqrt{3} - 3\sqrt{9}\sqrt{3} + 2\sqrt{25} \times \sqrt{3} \\
 &= 4\sqrt{3} - 9\sqrt{3} + 10\sqrt{3} = (4 - 9 + 10)\sqrt{3} = 5\sqrt{3}
 \end{aligned}$$

$$\begin{aligned}
 (ii) \quad &\sqrt[3]{128} - \sqrt[3]{250} + \sqrt[3]{432} \\
 &= \sqrt[3]{64 \times 2} - \sqrt[3]{125 \times 2} + \sqrt[3]{216 \times 2} \\
 &= \sqrt[3]{(4)^3 \times 2} - \sqrt[3]{(5)^3 \times 2} + \sqrt[3]{(6)^3 \times 2} \\
 &= \sqrt[3]{(4)^3} \sqrt[3]{2} - \sqrt[3]{(5)^3} \sqrt[3]{2} + \sqrt[3]{(6)^3} \sqrt[3]{2} \\
 &= 4\sqrt[3]{2} - 5\sqrt[3]{2} + 6\sqrt[3]{2} = (4 - 5 + 6)\sqrt[3]{2} = 5\sqrt[3]{2}
 \end{aligned}$$

### Example

Simplify and express the answer in the simplest form.

$$(i) \quad \sqrt{14}\sqrt{35}$$

$$(ii) \quad \frac{\sqrt[6]{12}}{\sqrt{3}\sqrt[3]{2}}$$

### Solution

$$(i) \quad \sqrt{14}\sqrt{35} = \sqrt{14 \times 35} = \sqrt{7 \times 2 \times 7 \times 5} = \sqrt{(7)^2 \times 2 \times 5}$$

$$= \sqrt{(7)^2 \times 10} = \sqrt{(7)^2} \times \sqrt{10} = 7\sqrt{10}$$

(ii) We have  $\frac{\sqrt[6]{12}}{\sqrt{3}\sqrt[3]{2}}$

For  $\sqrt{3} \sqrt[3]{2}$  the L.C.M of orders 2 and 3 is 6.

$$\text{Thus } \sqrt{3} = (3)^{1/2} = (3)^{3/6} = \sqrt[6]{3^3} = \sqrt[6]{27}$$

$$\text{and } \sqrt[3]{2} = (2)^{1/3} = (2)^{2/6} = \sqrt[6]{(2)^2} = \sqrt[6]{4}$$

$$\text{Hence } \frac{\sqrt[6]{12}}{\sqrt{3}\sqrt[3]{2}} = \frac{\sqrt[6]{12}}{\sqrt[6]{27}\sqrt[6]{4}} = \frac{\sqrt[6]{12}}{\sqrt[6]{108}} = \sqrt[6]{\frac{12}{108}} = \sqrt[6]{\frac{1}{9}}$$

Its simplest form is

$$\sqrt[6]{\left(\frac{1}{3}\right)^2} = \left(\frac{1}{3}\right)^{2/6} = \left(\frac{1}{3}\right)^{1/3} = \sqrt[3]{\frac{1}{3}}$$

### Exercise 4.3

1. Express each of the following surd in the simplest form.

(i)  $\sqrt{180}$

$$= \sqrt{2 \times 2 \times 3 \times 3 \times 5}$$

$$= 2 \times 3 \sqrt{5}$$

$$= 6\sqrt{5}$$

(ii)  $3\sqrt{162}$

$$= 3\sqrt{2 \times 3 \times 3 \times 3 \times 3}$$

$$= 3(3 \times 3)\sqrt{2}$$

$$= 27\sqrt{2}$$

(iii)  $\frac{3}{4}\sqrt[3]{128}$

$$= \frac{3}{4}(128)^{\frac{1}{3}}$$

$$= \frac{3}{4}(2 \times 2 \times 2 \times 2 \times 2 \times 2)^{\frac{1}{3}}$$

$$= \frac{3}{4}(2^3 \times 2^3 \times 2)^{\frac{1}{3}}$$

$$= \frac{3}{4}(2^3)^{\frac{1}{3}} \times (2^3)^{\frac{1}{3}} \times 2^{\frac{1}{3}}$$

$$= \frac{3}{4}(2)(2) \times \sqrt[3]{2}$$

$$= 3\sqrt[3]{2}$$

(iv)  $\sqrt[5]{96x^6y^7z^8}$

$$= \sqrt[5]{2 \times 2 \times 2 \times 2 \times 2 \times 3x^6y^7z^8}$$

$$= (2^5 \times 3x^5 \times y^5 \times z^5)^{\frac{1}{5}}$$

$$= (2^{\frac{1}{5}} \times 3^{\frac{1}{5}} \times x^{\frac{1}{5}} \times y^{\frac{1}{5}} \times z^{\frac{1}{5}})^{\frac{1}{5}}$$

$$= 2^{\frac{1}{5}} \times 3^{\frac{1}{5}} \times x^{\frac{1}{5}} \times y^{\frac{2}{5}} \times z^{\frac{3}{5}}$$

$$= 2xyz^{\frac{1}{5}} \times x^{\frac{1}{5}} \times y^{\frac{2}{5}} \times z^{\frac{3}{5}}$$

$$= 2xyz\sqrt[5]{3xy^2z^3}$$

**2. Simplify**

$$(i) \frac{\sqrt{18}}{\sqrt{3}\sqrt{2}} = \frac{\sqrt{3 \cdot 3 \cdot 2}}{\sqrt{3} \cdot \sqrt{2}} = \frac{3\sqrt{2}}{\sqrt{3}\sqrt{2}}$$

$$= \frac{3}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{3\sqrt{3}}{3} = \sqrt{3}$$

$$(ii) \frac{\sqrt{21} \times \sqrt{9}}{\sqrt{63}} = \frac{\sqrt{3 \times 7} \times \sqrt{3 \times 3}}{\sqrt{3 \times 3 \times 7}}$$

$$= \frac{\sqrt{3 \times 7 \times 3 \times 3}}{\sqrt{3 \times 3 \times 7}}$$

$$= \frac{3\sqrt{21}}{3\sqrt{7}} = \sqrt{\frac{21}{7}} = \sqrt{3}$$

$$(iii) \sqrt[5]{243x^5y^{10}z^{15}}$$

$$= (3^5 \cdot x^5 y^{10} z^{15})^{\frac{1}{5}}$$

$$= (3^5)^{\frac{1}{5}} (x^5)^{\frac{1}{5}} (y^{10})^{\frac{1}{5}} (z^{15})^{\frac{1}{5}}$$

$$= 3xy^2z^3$$

$$(iv) \frac{4}{5}\sqrt[3]{125}$$

$$= \frac{4}{5} \left( \cancel{5} \right)^{\frac{1}{3}} \cancel{5}$$

$$= 4$$

$$(v) \sqrt{21} \times \sqrt{7} \times \sqrt{3}$$

$$= \sqrt{3 \times 7} \times \sqrt{7} \times \sqrt{3}$$

$$= \sqrt{3 \times 7 \times 7 \times 3} = (3^2 \times 7^2)^{\frac{1}{2}}$$

$$= (3^2)^{\frac{1}{2}} x (7^2)^{\frac{1}{2}}$$

$$= 3 \times 7$$

$$= 21$$

**3. Simplify by combining similar terms:**

$$(i) \sqrt{45} - 3\sqrt{20} + 4\sqrt{5}$$

$$= \sqrt{9 \times 5} - 3\sqrt{4 \times 5} + 4\sqrt{5}$$

$$= 3\sqrt{5} - 6\sqrt{5} + 4\sqrt{5}$$

$$= (3 - 6 + 4)\sqrt{5}$$

$$= (-3 + 4)\sqrt{5}$$

$$= \sqrt{5}$$

$$(ii) 4\sqrt{12} + 5\sqrt{27} - 3\sqrt{75} + \sqrt{300}$$

$$= 4\sqrt{3 \times 4} + 5\sqrt{3 \times 3 \times 3} - 3\sqrt{3 \times 5 \times 5}$$

$$+ \sqrt{3 \times 2 \times 5 \times 2 \times 5}$$

$$= 8\sqrt{3} + 15\sqrt{3} - 15\sqrt{3} + 10\sqrt{3}$$

$$= (8 + 15 - 15 + 10)\sqrt{3}$$

$$= 18\sqrt{3}$$

$$(iii) \sqrt{3}(2\sqrt{3} + 3\sqrt{3})$$

$$= \sqrt{3}((2+3)\sqrt{3})$$

$$= \sqrt{3}(5\sqrt{3})$$

$$= 5\sqrt{3} \times \sqrt{3}$$

$$= 5(\sqrt{3 \times 3})$$

$$= 5(3)$$

$$= 15$$

$$(iv) 2(6\sqrt{5} - 3\sqrt{5})$$

$$= 2((6-3)\sqrt{5})$$

$$= 2(3\sqrt{5})$$

$$= 6\sqrt{5}$$

**4. Simplify:**

$$(i) (3 + \sqrt{3})(3 - \sqrt{3})$$

$$= (3)^2 - (\sqrt{3})^2$$

$$\begin{aligned}
 &= 9 - 3 \\
 &= 6 \\
 \text{(ii)} \quad &(\sqrt{5} + \sqrt{3})^2 \\
 &= (\sqrt{5})^2 + (\sqrt{3})^2 + 2\sqrt{5}\sqrt{3} \\
 &= 5 + 3 + 2\sqrt{15} \\
 &= 8 + 2\sqrt{15} \\
 \text{(iii)} \quad &(\sqrt{5} + \sqrt{3})(\sqrt{5} - \sqrt{3}) \\
 &= (\sqrt{5})^2 - (\sqrt{3})^2 \\
 &= 5 - 3 \\
 &= 2 \\
 \text{(iv)} \quad &\left(\sqrt{2} + \frac{1}{\sqrt{3}}\right)\left(\sqrt{2} - \frac{1}{\sqrt{3}}\right) \\
 &= (\sqrt{2})^2 - \left(\frac{1}{\sqrt{3}}\right)^2
 \end{aligned}$$

$$\begin{aligned}
 &= 2 - \frac{1}{3} \\
 &= \frac{6-1}{3} = \frac{5}{3} \\
 \text{(v)} \quad &(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y})(x + y) \\
 &\quad (x^2 + y^2) \\
 &= ((\sqrt{x})^2 - (\sqrt{y})^2)(x + y)(x^2 + y^2) \\
 &= (x - y)(x + y)(x^2 + y^2) \\
 &= (x^2 - y^2)(x^2 + y^2) \\
 &= (x^2)^2 - (y^2)^2 \\
 &= x^4 - y^4
 \end{aligned}$$

### Define monomial surd

- (i) A surd which contains a single term is called a monomial surd. e.g.,  $\sqrt{2}, \sqrt{3}$  etc.
- (ii) A surd which contains sum of two monomial surds or sum of a monomial surd and a rational number is called a binomial surd.  
e.g.,  $\sqrt{3} + \sqrt{7}$  or  $\sqrt{2} + 5\sqrt{11} - 8$  etc.
- (iii) If the product of two surds is a rational number, then each surd is called the rationalizing factor of the other.

### Example

Rationalize the denominator  $\frac{58}{7-2\sqrt{5}}$

### Solution

To rationalize the denominator, we multiply both the numerator and denominator by the conjugate  $(7+2\sqrt{5})$  of  $(7-2\sqrt{5})$ , i.e.

$$\frac{58}{7-2\sqrt{5}} = \frac{58}{7-2\sqrt{5}} \times \frac{7+2\sqrt{5}}{7+2\sqrt{5}} = \frac{58(7+2\sqrt{5})}{(7)^2 - (2\sqrt{5})^2}$$

$$\begin{aligned}
 &= \frac{58(7+2\sqrt{5})}{49-20}; \text{ (radical is eliminated in the denominator)} \\
 &= \frac{58(7+2\sqrt{5})}{29} = 2(7+2\sqrt{5})
 \end{aligned}$$

### Example

Rationalize the denominator  $\frac{2}{\sqrt{5}+\sqrt{2}}$

### Solution

Multiplying both the numerator and denominator by the conjugate  $(\sqrt{5}-\sqrt{2})$  of  $(\sqrt{5}+\sqrt{2})$ , to get

$$\begin{aligned}
 \frac{2}{\sqrt{5}+\sqrt{2}} &= \frac{2}{\sqrt{5}+\sqrt{2}} \times \frac{\sqrt{5}-\sqrt{2}}{\sqrt{5}-\sqrt{2}} = \frac{2(\sqrt{5}-\sqrt{2})}{(\sqrt{5})^2-(\sqrt{2})^2} = \frac{2(\sqrt{5}-\sqrt{2})}{5-2} \\
 &= \frac{2(\sqrt{5}-\sqrt{2})}{3} = \frac{2(\sqrt{5}-\sqrt{2})}{3}
 \end{aligned}$$

### Example

Simplify  $\frac{6}{2\sqrt{3}-\sqrt{6}} + \frac{\sqrt{6}}{\sqrt{3}+\sqrt{2}} - \frac{4\sqrt{3}}{\sqrt{6}-\sqrt{2}}$

### Solution

First we shall rationalize the denominators and then simplify. We have

$$\begin{aligned}
 &\frac{6}{2\sqrt{3}-\sqrt{6}} + \frac{\sqrt{6}}{\sqrt{3}+\sqrt{2}} - \frac{4\sqrt{3}}{\sqrt{6}-\sqrt{2}} \\
 &= \frac{6}{2\sqrt{3}-\sqrt{6}} \times \frac{2\sqrt{3}+\sqrt{6}}{2\sqrt{3}+\sqrt{6}} + \frac{\sqrt{6}}{\sqrt{3}+\sqrt{2}} \times \frac{\sqrt{3}-\sqrt{2}}{\sqrt{3}-\sqrt{2}} - \frac{4\sqrt{3}}{\sqrt{6}-\sqrt{2}} \times \frac{\sqrt{6}+\sqrt{2}}{\sqrt{6}+\sqrt{2}} \\
 &= \frac{6(2\sqrt{3}+\sqrt{6})}{(2\sqrt{3})^2-(\sqrt{6})^2} + \frac{\sqrt{6}(\sqrt{3}-\sqrt{2})}{(\sqrt{3})^2-(\sqrt{2})^2} - \frac{4\sqrt{3}(\sqrt{6}+\sqrt{2})}{(\sqrt{6})^2-(\sqrt{2})^2} \\
 &= \frac{6(2\sqrt{3}+\sqrt{6})}{12-6} + \frac{\sqrt{6}(\sqrt{3}-\sqrt{2})}{3-2} - \frac{4\sqrt{3}(\sqrt{6}+\sqrt{2})}{6-2} \\
 &= \frac{12\sqrt{3}+6\sqrt{6}}{6} + \frac{\sqrt{6}\sqrt{3}-\sqrt{6}\sqrt{2}}{1} - \frac{4\sqrt{3}\sqrt{6}+4\sqrt{3}\sqrt{2}}{4} \\
 &= 2\sqrt{3}+\sqrt{6}+3\sqrt{2}-2\sqrt{3}-3\sqrt{2}-\sqrt{6} = 0
 \end{aligned}$$

**Example**

Find rational numbers  $x$  and  $y$  such that  $\frac{4+3\sqrt{5}}{4-3\sqrt{5}} = x + y\sqrt{5}$

**Solution**

$$\begin{aligned}\frac{4+3\sqrt{5}}{4-3\sqrt{5}} &= \frac{4+3\sqrt{5}}{4-3\sqrt{5}} \times \frac{4+3\sqrt{5}}{4+3\sqrt{5}} = \frac{(4+3\sqrt{5})^2}{(4)^2 - (3\sqrt{5})^2} \\ &= \frac{16+24\sqrt{5}+45}{16-45} = \frac{61+24\sqrt{5}}{-29} \\ \Rightarrow \frac{-61}{29} - \frac{24}{29}\sqrt{5} &= x + y\sqrt{5} \quad (\text{given})\end{aligned}$$

Hence, on comparing the two sides, we get

$$x = \frac{-61}{29}, \quad y = \frac{-24}{29}$$

**Example**

If  $x = 3 + \sqrt{8}$ , then evaluate

$$(i) \quad x + \frac{1}{x} \quad \text{and} \quad (ii) \quad x^2 + \frac{1}{x^2}$$

**Solution**

Since  $x = 3 + \sqrt{8}$ , therefore,

$$\begin{aligned}\frac{1}{x} &= \frac{1}{3+\sqrt{8}} = \frac{1}{3+\sqrt{8}} \times \frac{3-\sqrt{8}}{3-\sqrt{8}} = \frac{3-\sqrt{8}}{(3)^2 - (\sqrt{8})^2} \\ &= \frac{3-\sqrt{8}}{9-8} = 3-\sqrt{8}\end{aligned}$$

$$(i) \quad x + \frac{1}{x} = 3 + \sqrt{8} + 3 - \sqrt{8} = 6$$

$$(ii) \quad \left(x + \frac{1}{x}\right)^2 = 36$$

$$\text{or} \quad x^2 + 2x \times \frac{1}{x} + \frac{1}{x^2} = 36$$

$$\text{or} \quad x^2 + \frac{1}{x^2} = 34$$

## Exercise 4.4

**1. Rationalize the denominator**

$$\begin{aligned} \text{(i)} \quad \frac{3}{4\sqrt{3}} &= \frac{3}{4\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{3\sqrt{3}}{4\sqrt{3} \times 3} \\ &= \frac{3\sqrt{3}}{4(3)} = \frac{\sqrt{3}}{4} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \frac{14}{\sqrt{98}} &= \frac{14}{7\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} \\ &= \frac{14\sqrt{2}}{14} = \sqrt{2} \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad \frac{6}{\sqrt{8} \cdot \sqrt{27}} &= \frac{6}{2\sqrt{2} \cdot 3\sqrt{3}} \\ &= \frac{6}{6\sqrt{6}} \\ &= \frac{1}{\sqrt{6}} \\ &= \frac{1}{\sqrt{6}} \times \frac{\sqrt{6}}{\sqrt{6}} \\ &= \frac{\sqrt{6}}{6} \end{aligned}$$

$$\begin{aligned} \text{(iv)} \quad \frac{1}{3+2\sqrt{5}} &= \frac{1}{3+2\sqrt{5}} \times \frac{3-2\sqrt{5}}{3-2\sqrt{5}} \\ &= \frac{3-2\sqrt{5}}{(3)^2 - (2\sqrt{5})^2} = \frac{3-2\sqrt{5}}{9-20} \\ &= \frac{3-2\sqrt{5}}{-11} \end{aligned}$$

$$\begin{aligned} \text{(v)} \quad \frac{15}{\sqrt{31}-4} &= \frac{15}{\sqrt{31}-4} \times \frac{\sqrt{31}+4}{\sqrt{31}+4} \end{aligned}$$

$$= \frac{15(\sqrt{31}+4)}{(\sqrt{31})^2 - (4)^2}$$

$$= \frac{15(\sqrt{31}+4)}{31-16}$$

$$= \frac{15(\sqrt{31}+4)}{15}$$

$$= \sqrt{31} + 4$$

$$\begin{aligned} \text{(vi)} \quad \frac{2}{\sqrt{5}-\sqrt{3}} &= \frac{2}{\sqrt{5}-\sqrt{3}} \times \frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}+\sqrt{3}} \\ &= \frac{2(\sqrt{5}+\sqrt{3})}{(\sqrt{5})^2 - (\sqrt{3})^2} \\ &= \frac{2(\sqrt{5}+\sqrt{3})}{5-3} \\ &= \frac{2(\sqrt{5}+\sqrt{3})}{2} \\ &= \sqrt{5} + \sqrt{3} \end{aligned}$$

$$\begin{aligned} \text{(vii)} \quad \frac{\sqrt{3}-1}{\sqrt{3}+1} &= \frac{\sqrt{3}-1}{\sqrt{3}+1} \times \frac{\sqrt{3}-1}{\sqrt{3}-1} \\ &= \frac{(\sqrt{3}-1)(\sqrt{3}-1)}{(\sqrt{3})^2 - (1)^2} \end{aligned}$$

$$= \frac{(\sqrt{3}-1)^2}{3-1}$$

$$\begin{aligned} &= \frac{(\sqrt{3})^2 + 1^2 - 2(1)\sqrt{3}}{2} \\ &= \frac{3+1-2\sqrt{3}}{2} \end{aligned}$$

$$\begin{aligned}
 &= \frac{4-2\sqrt{3}}{2} \\
 &= \frac{2(2-\sqrt{3})}{2} \\
 &= 2-\sqrt{3} \\
 (\text{viii}) \quad &\frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}-\sqrt{3}} = \frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}-\sqrt{3}} \times \frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}+\sqrt{3}} \\
 &= \frac{(\sqrt{5}+\sqrt{3})^2}{(\sqrt{5})^2 - (\sqrt{3})^2} \\
 &= \frac{(\sqrt{5}+\sqrt{3})^2}{5-3} \\
 &= \frac{(\sqrt{5})^2 + (\sqrt{3})^2 + 2(\sqrt{5})(\sqrt{3})}{2} \\
 &= \frac{5+3+2\sqrt{15}}{2} \\
 &= \frac{8+2\sqrt{15}}{2} \\
 &= \frac{2(4+\sqrt{15})}{2} \\
 &= 4+\sqrt{15}
 \end{aligned}$$

(2) Find conjugate of  $x+\sqrt{y}$ :

- (i)  $3+\sqrt{7}$   
Conjugate of  $3+\sqrt{7}$  is  $3-\sqrt{7}$
- (ii)  $4-\sqrt{5}$   
Conjugate of  $4-\sqrt{5}$  is  $4+\sqrt{5}$
- (iii)  $2+\sqrt{3}$   
Conjugate of  $2+\sqrt{3}$  is  $2-\sqrt{3}$
- (iv)  $2+\sqrt{5}$   
Conjugate of  $2+\sqrt{5}$  is  $2-\sqrt{5}$
- (v)  $5+\sqrt{7}$

Conjugate of  $5+\sqrt{7}$  is  $5-\sqrt{7}$

- (vi)  $4-\sqrt{15}$   
Conjugate of  $4-\sqrt{15}$  is  $4+\sqrt{15}$
- (vii)  $7-\sqrt{6}$   
Conjugate of  $7-\sqrt{6}$  is  $7+\sqrt{6}$
- (viii)  $9+\sqrt{2}$   
Conjugate of  $9+\sqrt{2}$  is  $9-\sqrt{2}$

Q.3 If  $x=2-\sqrt{3}$  find  $\frac{1}{x}$

$$\begin{aligned}
 \text{(i)} \quad x &= 2-\sqrt{3} \\
 \frac{1}{x} &= \frac{1}{2-\sqrt{3}} \times \frac{2+\sqrt{3}}{2+\sqrt{3}} \\
 \frac{1}{x} &= \frac{2+\sqrt{3}}{(2)^2 - (\sqrt{3})^2} \\
 \frac{1}{x} &= \frac{2+\sqrt{3}}{4-3} \\
 \frac{1}{x} &= 2+\sqrt{3} \\
 \text{(ii)} \quad x=4-\sqrt{17} \quad \text{find } \frac{1}{x} \\
 \frac{1}{x} &= \frac{1}{4-\sqrt{17}} \times \frac{4+\sqrt{17}}{4+\sqrt{17}} \\
 \frac{1}{x} &= \frac{4+\sqrt{17}}{(4)^2 - (\sqrt{17})^2} \\
 &= \frac{4+\sqrt{17}}{16-17} \\
 &= \frac{4+\sqrt{17}}{-1} \\
 &= -(4+\sqrt{17}) \\
 &= -4-\sqrt{17}
 \end{aligned}$$

(iii) If  $x = \sqrt{3} + 2$ , find  $x + \frac{1}{x}$

$$x = \sqrt{3} + 2$$

$$\frac{1}{x} = \frac{1}{\sqrt{3} + 2} \times \frac{\sqrt{3} - 2}{\sqrt{3} - 2}$$

$$\frac{1}{x} = \frac{\sqrt{3} - 2}{(\sqrt{3})^2 - (2)^2}$$

$$\frac{1}{x} = \frac{\sqrt{3} - 2}{3 - 4}$$

$$\frac{1}{x} = \frac{\sqrt{3} - 2}{-1}$$

$$\frac{1}{x} = -\sqrt{3} + 2 = 2 - \sqrt{3}$$

$$x + \frac{1}{x} = \sqrt{3} + 2 - \sqrt{3} + 2$$

$$x + \frac{1}{x} = 4$$

#### Q4. Simplify

$$(i) \quad \frac{1+\sqrt{2}}{\sqrt{5}+\sqrt{3}} + \frac{1-\sqrt{2}}{\sqrt{5}-\sqrt{3}}$$

$$\begin{aligned} & \frac{1+\sqrt{2}}{\sqrt{5}+\sqrt{3}} \times \frac{\sqrt{5}-\sqrt{3}}{\sqrt{5}-\sqrt{3}} + \frac{1-\sqrt{2}}{\sqrt{5}-\sqrt{3}} \times \frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}+\sqrt{3}} \\ &= \frac{(1+\sqrt{2})(\sqrt{5}-\sqrt{3})}{(\sqrt{5})^2 - (\sqrt{3})^2} + \frac{(1-\sqrt{2})(\sqrt{5}+\sqrt{3})}{(\sqrt{5})^2 - (\sqrt{3})^2} \end{aligned}$$

$$\begin{aligned} &= \frac{(1+\sqrt{2})(\sqrt{5}-\sqrt{3})}{5-3} + \frac{(1-\sqrt{2})(\sqrt{5}+\sqrt{3})}{5-3} \\ &= \frac{\sqrt{5}-\sqrt{3}+\sqrt{2}\sqrt{5}-\sqrt{2}\sqrt{3}}{2} + \frac{\sqrt{5}+\sqrt{3}-\sqrt{2}\sqrt{5}-\sqrt{2}\sqrt{3}}{2} \\ &= \frac{\sqrt{5}-\sqrt{3}+\sqrt{10}-\sqrt{6}}{2} + \frac{\sqrt{5}+\sqrt{3}-\sqrt{10}-\sqrt{6}}{2} \end{aligned}$$

$$= \frac{\sqrt{5}-\sqrt{3}+\sqrt{10}-\sqrt{6}+\sqrt{5}+\sqrt{3}-\sqrt{10}-\sqrt{6}}{2}$$

$$= \frac{2\sqrt{5}-2\sqrt{6}}{2}$$

$$= \frac{\cancel{2}(\sqrt{5}-\sqrt{6})}{\cancel{2}}$$

$$= \sqrt{5}-\sqrt{6}$$

$$(ii) \quad \frac{1}{2+\sqrt{3}} + \frac{2}{\sqrt{5}-\sqrt{3}} + \frac{1}{2+\sqrt{5}}$$

$$\begin{aligned} &= \frac{1}{2+\sqrt{3}} \times \frac{2-\sqrt{3}}{2-\sqrt{3}} + \frac{2}{\sqrt{5}-\sqrt{3}} \\ &\quad \times \frac{\sqrt{5}+\sqrt{3}}{\sqrt{5}+\sqrt{3}} + \frac{1}{2+\sqrt{5}} \times \frac{2-\sqrt{5}}{2-\sqrt{5}} \end{aligned}$$

$$= \frac{2-\sqrt{3}}{(2)^2 - (\sqrt{3})^2} + \frac{2(\sqrt{5}+\sqrt{3})}{(\sqrt{5})^2 - (\sqrt{3})^2} + \frac{2-\sqrt{5}}{(2)^2 - (\sqrt{5})^2}$$

$$= \frac{2-\sqrt{3}}{4-3} + \frac{2(\sqrt{5}+\sqrt{3})}{5-3} + \frac{2-\sqrt{5}}{4-5}$$

$$= 2-\sqrt{3} + \frac{2(\sqrt{5}+\sqrt{3})}{2} + \frac{2-\sqrt{5}}{-1}$$

$$= \cancel{2} - \cancel{\sqrt{3}} + \sqrt{5} + \sqrt{3} - \cancel{2} + \sqrt{5} = 2\sqrt{5}$$

$$(iii) \quad \frac{2}{\sqrt{5}+\sqrt{3}} + \frac{1}{\sqrt{3}+\sqrt{2}} - \frac{3}{\sqrt{5}+\sqrt{2}}$$

$$= \frac{2}{\sqrt{5}+\sqrt{3}} \times \frac{\sqrt{5}-\sqrt{3}}{\sqrt{5}-\sqrt{3}} + \frac{1}{\sqrt{3}+\sqrt{2}}$$

$$\times \frac{\sqrt{3}-\sqrt{2}}{\sqrt{3}-\sqrt{2}} - \frac{3}{\sqrt{5}+\sqrt{2}} \times \frac{\sqrt{5}-\sqrt{2}}{\sqrt{5}-\sqrt{2}}$$

$$= \frac{2(\sqrt{5}-\sqrt{3})}{(\sqrt{5})^2 - (\sqrt{3})^2} + \frac{\sqrt{3}-\sqrt{2}}{(\sqrt{3})^2 - (\sqrt{2})^2} - \frac{3(\sqrt{5}-\sqrt{2})}{(\sqrt{5})^2 - (\sqrt{2})^2}$$

$$= \frac{2(\sqrt{5}-\sqrt{3})}{5-3} + \frac{\sqrt{3}-\sqrt{2}}{3-2} - \frac{3(\sqrt{5}-\sqrt{2})}{5-2}$$

$$\begin{aligned}
 &= \frac{\cancel{2}(\sqrt{5}-\sqrt{3})}{\cancel{2}} + \frac{\sqrt{3}-\sqrt{2}}{1} - \frac{\cancel{2}(\sqrt{5}-\sqrt{2})}{\cancel{2}} \\
 &= \cancel{\sqrt{5}} - \cancel{\sqrt{3}} + \cancel{\sqrt{3}} - \cancel{\sqrt{2}} - \cancel{\sqrt{5}} + \cancel{\sqrt{2}} \\
 &= 0
 \end{aligned}$$

**Q5(i)** If  $x = 2 + \sqrt{3}$ , find value of  $x - \frac{1}{x}$

and  $\left(x - \frac{1}{x}\right)^2$

$$x = 2 + \sqrt{3}$$

$$\frac{1}{x} = \frac{1}{2+\sqrt{3}} \times \frac{2-\sqrt{3}}{2-\sqrt{3}}$$

$$\frac{1}{x} = \frac{2-\sqrt{3}}{(2)^2 - (\sqrt{3})^2}$$

$$\frac{1}{x} = 2 - \sqrt{3}$$

$$x - \frac{1}{x} = 2 + \sqrt{3} - (2 - \sqrt{3})$$

$$\begin{aligned}
 &= \cancel{2} + \sqrt{3} - \cancel{2} + \sqrt{3} \\
 &= 2\sqrt{3}
 \end{aligned}$$

$$\left(x - \frac{1}{x}\right)^2 = (2\sqrt{3})^2$$

$$\left(x - \frac{1}{x}\right)^2 = 12$$

**(ii)** If  $x = \frac{\sqrt{5}-\sqrt{2}}{\sqrt{5}+\sqrt{2}}$  find the value of

$$x + \frac{1}{x}, x^2 + \frac{1}{x^2} \text{ and } x^3 + \frac{1}{x^3}$$

$$x = \frac{\sqrt{5}-\sqrt{2}}{\sqrt{5}+\sqrt{2}} \times \frac{\sqrt{5}-\sqrt{2}}{\sqrt{5}-\sqrt{2}}$$

$$x = \frac{(\sqrt{5}-\sqrt{2})^2}{(\sqrt{5})^2 - (\sqrt{2})^2}$$

$$x = \frac{(\sqrt{5})^2 + (\sqrt{2})^2 - 2(\sqrt{5})(\sqrt{2})}{5-2}$$

$$x = \frac{5+2-2\sqrt{10}}{3}$$

$$x = \frac{7-2\sqrt{10}}{3}$$

$$\frac{1}{x} = \frac{3}{7-2\sqrt{10}} \times \frac{7+2\sqrt{10}}{7+2\sqrt{10}}$$

$$\frac{1}{x} = \frac{3(7+2\sqrt{10})}{(7)^2 - (2\sqrt{10})^2}$$

$$\frac{1}{x} = \frac{3(7+2\sqrt{10})}{49-40}$$

$$\frac{1}{x} = \frac{3(7+2\sqrt{10})}{9}$$

$$\frac{1}{x} = \frac{7+2\sqrt{10}}{3}$$

$$\begin{aligned}
 x + \frac{1}{x} &= \frac{7-2\sqrt{10}}{3} + \frac{7+2\sqrt{10}}{3} \\
 &= \frac{7-2\sqrt{10} + 7+2\sqrt{10}}{3} = \frac{14}{3}
 \end{aligned}$$

Now

$$x + \frac{1}{x} = \frac{14}{3}$$

Squaring

$$\left(x + \frac{1}{x}\right)^2 = \left(\frac{14}{3}\right)^2$$

$$x^2 + \frac{1}{x^2} + 2 = \frac{196}{9}$$

$$x^2 + \frac{1}{x^2} = \frac{196}{9} - 2$$

$$x^2 + \frac{1}{x^2} = \frac{196-18}{9} = \frac{178}{9}$$

Also

$$x^3 + \frac{1}{x^3} = ?$$

$$x + \frac{1}{x} = \frac{14}{3}$$

$$\left( x + \frac{1}{x} \right)^3 = \left( \frac{14}{3} \right)^3$$

$$x^3 + \frac{1}{x^3} + 3(x) \left( \frac{1}{x} \right) \left( x + \frac{1}{x} \right) = \frac{2744}{27}$$

$$x^3 + \frac{1}{x^3} + 3 \left( x + \frac{1}{x} \right) = \frac{2744}{27}$$

$$x^3 + \frac{1}{x^3} + 3 \left( \frac{14}{3} \right) = \frac{2744}{27}$$

$$\begin{aligned} x^3 + \frac{1}{x^3} &= \frac{2744}{27} - 14 \\ &= \frac{2366}{27} \end{aligned}$$

**Q6.** Determine the rational numbers a and b. If

$$\frac{\sqrt{3}-1}{\sqrt{3}+1} + \frac{\sqrt{3}+1}{\sqrt{3}-1} = a+b\sqrt{3}$$

**Given**

**Q1.** If  $x + \frac{1}{x} = 3$  find

$$(i) \quad x^2 + \frac{1}{x^2}$$

$$(ii) \quad x^4 + \frac{1}{x^4}$$

$$(i) \quad x + \frac{1}{x} = 3$$

$$\left( x + \frac{1}{x} \right)^2 = (3)^2$$

$$x^2 + \frac{1}{x^2} + 2 = 9$$

$$x^2 + \frac{1}{x^2} = 9 - 2$$

$$\frac{\sqrt{3}-1}{\sqrt{3}+1} + \frac{\sqrt{3}+1}{\sqrt{3}-1} = a+b\sqrt{3}$$

$$\frac{\sqrt{3}-1}{\sqrt{3}+1} \times \frac{\sqrt{3}-1}{\sqrt{3}-1} + \frac{\sqrt{3}+1}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1} = a+b\sqrt{3}$$

$$\frac{(\sqrt{3}-1)^2}{(\sqrt{3})^2 - (1)^2} + \frac{(\sqrt{3}+1)^2}{(\sqrt{3})^2 - (1)^2} = a+b\sqrt{3}$$

$$\frac{(\sqrt{3})^2 + (1)^2 - 2(\sqrt{3})(1)}{3-1} + \frac{(\sqrt{3})^2 + (1)^2 + 2\sqrt{3}}{3-1} = a+b\sqrt{3}$$

$$\frac{3+1-2\sqrt{3}}{2} + \frac{3+1+2\sqrt{3}}{2} = a+b\sqrt{3}$$

$$\frac{4-2\sqrt{3}}{2} + \frac{4+2\sqrt{3}}{2} = a+b\sqrt{3}$$

$$\cancel{2}(2-\sqrt{3}) + \cancel{2}(2+\sqrt{3}) = a+b\sqrt{3}$$

$$2-\cancel{2}\sqrt{3} + 2 + \cancel{2}\sqrt{3} = a+b\sqrt{3}$$

$$4 = a+b\sqrt{3}$$

$$\Rightarrow a+b\sqrt{3} = 4$$

Hence on comparing the two sides, we get

$$\Rightarrow a=4 \text{ and } b=0$$

## Exercise

$$x^2 + \frac{1}{x^2} = 7$$

$$(iii) \quad x^4 + \frac{1}{x^4}$$

$$\left( x^2 + \frac{1}{x^2} \right)^2 = (7)^2$$

$$x^4 + \frac{1}{x^4} + 2 = 49$$

$$x^4 + \frac{1}{x^4} = 49 - 2$$

$$x^4 + \frac{1}{x^4} = 47$$

**Q2.** If  $x - \frac{1}{x} = 2$  find

$$(i) x^2 + \frac{1}{x^2}$$

$$(i) x - \frac{1}{x} = 2$$

Squaring

$$\left(x - \frac{1}{x}\right)^2 = (2)^2$$

$$x^2 + \frac{1}{x^2} - 2 = 4$$

$$x^2 + \frac{1}{x^2} = 4 + 2$$

$$x^2 + \frac{1}{x^2} = 6$$

$$(ii) \left(x^2 + \frac{1}{x^2}\right) = (6)^2$$

$$x^4 + \frac{1}{x^4} + 2 = 36$$

$$x^4 + \frac{1}{x^4} = 36 - 2$$

$$x^4 + \frac{1}{x^4} = 34$$

**Q3.** Find value of  $x^3 + y^3$  and  $xy$  if

$$x + y = 5 \text{ and } x - y = 3$$

$$4xy = (x + y)^2 - (x - y)^2$$

$$4xy = (5)^2 - (3)^2$$

Now

$$4xy = 25 - 9 = 16$$

$$xy = \frac{16}{4} = 4$$

$$x + y = 5$$

taking cube both sides

$$(x + y)^3 = (5)^3$$

$$x^3 + y^3 + 3xy(x + y) = 125$$

$$x^3 + y^3 + 3(4)(5) = 125$$

$$x^3 + y^3 + 60 = 125$$

$$x^3 + y^3 = 125 - 60$$

$$x^3 + y^3 = 65$$

**Q4.** If  $P = 2 + \sqrt{3}$  find (i)  $P + \frac{1}{P}$

$$(ii) P - \frac{1}{P} \quad (iii) P^2 + \frac{1}{P^2} \quad (iv) P^2 - \frac{1}{P^2}$$

$$P = 2 + \sqrt{3}$$

$$\frac{1}{P} = \frac{1}{2 + \sqrt{3}} \times \frac{2 - \sqrt{3}}{2 - \sqrt{3}}$$

$$\frac{1}{P} = \frac{2 - \sqrt{3}}{(2)^2 - (\sqrt{3})^2} = \frac{2 - \sqrt{3}}{4 - 3} = 2 - \sqrt{3}$$

$$i) P + \frac{1}{P} = 2 + \sqrt{3} + 2 - \sqrt{3} = 4$$

$$ii) P - \frac{1}{P} = 2 + \sqrt{3} - 2 - \sqrt{3} = 2\sqrt{3}$$

$$iii) P^2 + \frac{1}{P^2} = ?$$

$$\left(P + \frac{1}{P}\right)^2 = (4)^2$$

$$P^2 + \frac{1}{P^2} + 2 = 16$$

$$P^2 + \frac{1}{P^2} = 16 - 2$$

$$P^2 + \frac{1}{P^2} = 14$$

$$iv) P^2 - \frac{1}{P^2} = ?$$

$$\begin{aligned} P^2 - \frac{1}{P^2} &= \left( P + \frac{1}{P} \right) \left( P - \frac{1}{P} \right) \\ &= (4)(\sqrt{3}) \\ &= 8\sqrt{3} \end{aligned}$$

Q5. If  $q = \sqrt{5} + 2$  Find (i)  $q + \frac{1}{q}$

(ii)  $q - \frac{1}{q}$  (iii)  $q^2 + \frac{1}{q^2}$  (iv)  $q^2 - \frac{1}{q^2}$

**Solution:**  $q = \sqrt{5} + 2$

$$\frac{1}{q} = \frac{1}{\sqrt{5}+2} \times \frac{\sqrt{5}-2}{\sqrt{5}-2}$$

$$\frac{1}{q} = \frac{\sqrt{5}-2}{(\sqrt{5})^2 - (2)^2}$$

$$\frac{1}{q} = \frac{\sqrt{5}-2}{1} = \sqrt{5} - 2$$

(i)  $q + \frac{1}{q} = \sqrt{5} + 2 + \sqrt{5} - 2$   
 $= 2\sqrt{5}$

(ii)  $q - \frac{1}{q} = \sqrt{5} + 2 - \sqrt{5} + 2$   
 $= 4$

(iii)  $q^2 + \frac{1}{q^2}$   
 $\left( q + \frac{1}{q} \right)^2 = (2\sqrt{5})^2$

$$q^2 + \frac{1}{q^2} + 2 = 20$$

$$q^2 + \frac{1}{q^2} = 20 - 2$$

$$q^2 + \frac{1}{q^2} = 18$$

(iv)  $q^2 - \frac{1}{q^2} = \left( q + \frac{1}{q} \right) \left( q - \frac{1}{q} \right)$

$$\begin{aligned} &= (2\sqrt{5})(4) \\ &= 8\sqrt{5} \end{aligned}$$

### Q6. Simplify

$$\begin{aligned} \text{i)} \quad &\frac{\sqrt{a^2+2} + \sqrt{a^2-2}}{\sqrt{a^2+2} - \sqrt{a^2-2}} \\ &= \frac{\sqrt{a^2+2} + \sqrt{a^2-2}}{\sqrt{a^2+2} - \sqrt{a^2-2}} \times \frac{\sqrt{a^2+2} + \sqrt{a^2-2}}{\sqrt{a^2+2} + \sqrt{a^2-2}} \\ &= \frac{(\sqrt{a^2+2} + \sqrt{a^2-2})^2}{(\sqrt{a^2+2})^2 - (\sqrt{a^2-2})^2} \\ &= \frac{(\sqrt{a^2+2})^2 + (\sqrt{a^2-2})^2 + 2(\sqrt{a^2+2})(\sqrt{a^2-2})}{a^2 + 2 - a^2 + 2} \\ &= \frac{a^2 + 2 + a^2 - 2 + 2\sqrt{a^4 - 4}}{4} \\ &= \frac{2a^2 + 2\sqrt{a^4 - 4}}{4} \\ &= \frac{2(a^2 + \sqrt{a^4 - 4})}{4} \\ &= \frac{a^2 + \sqrt{a^4 - 4}}{2} \end{aligned}$$

$$\begin{aligned} \text{ii)} \quad &\frac{1}{a - \sqrt{a^2 - x^2}} - \frac{1}{a + \sqrt{a^2 - x^2}} \\ &= \frac{1}{a - \sqrt{a^2 - x^2}} \times \frac{a + \sqrt{a^2 - x^2}}{a + \sqrt{a^2 - x^2}} \\ &\quad - \frac{1}{a + \sqrt{a^2 - x^2}} \times \frac{a - \sqrt{a^2 - x^2}}{a - \sqrt{a^2 - x^2}} \\ &= \frac{a + \sqrt{a^2 - x^2}}{(a)^2 - (\sqrt{a^2 - x^2})^2} - \frac{a - \sqrt{a^2 - x^2}}{(a)^2 - (\sqrt{a^2 - x^2})^2} \end{aligned}$$

$$= \frac{a + \sqrt{a^2 - x^2}}{a^2 - a^2 + x^2} - \frac{a - \sqrt{a^2 - x^2}}{a^2 - a^2 + x^2}$$

$$= \frac{a + \sqrt{a^2 - x^2}}{x^2} - \frac{a - \sqrt{a^2 - x^2}}{x^2}$$

$$= \frac{a + \sqrt{a^2 - x^2} - a + \sqrt{a^2 - x^2}}{x^2}$$

$$= \frac{2\sqrt{a^2 - x^2}}{x^2}$$

## Objective

1.  $4x + 3y - 2$  is an algebraic \_\_\_\_\_  
 (a) Expression  
 (b) Sentence  
 (c) Equation  
 (d) In equation
2. The degree of polynomial  $4x^4 + 2x^2y$  is \_\_\_\_\_  
 (a) 1      (b) 2  
 (c) 3      (d) 4
3.  $a^3 + b^3$  is equal to \_\_\_\_\_  
 (a)  $(a-b)(a^2+ab+b^2)$   
 (b)  $(a+b)(a^2-ab+b^2)$   
 (c)  $(a-b)(a^2-ab+b^2)$   
 (d)  $(a-b)(a^2+ab-b^2)$
4.  $(3 + \sqrt{2})(3 - \sqrt{2})$  is equal to: \_\_\_\_\_  
 (a) 7      (b) -7  
 (c) -1      (d) 1
5. Conjugate of Surd  $a + \sqrt{b}$  is \_\_\_\_\_  
 (a)  $-a + \sqrt{b}$       (b)  $a - \sqrt{b}$   
 (d)  $\sqrt{a} + \sqrt{b}$       (d)  $\sqrt{a} - \sqrt{b}$
6.  $\frac{1}{a-b} - \frac{1}{a+b}$  is equal to  
 (a)  $\frac{2a}{a^2-b^2}$       (b)  $\frac{2b}{a^2-b^2}$   
 (c)  $\frac{-2a}{a^2-b^2}$       (d)  $\frac{-2b}{a^2-b^2}$

7.  $\frac{a^2 - b^2}{a+b}$  is equal to:  
 (a)  $(a-b)^2$       (b)  $(a+b)^2$   
 (c)  $a+b$       (d)  $a-b$
8.  $(\sqrt{a} + \sqrt{b})(\sqrt{a} - \sqrt{b})$  is equal to: \_\_\_\_\_  
 (a)  $a^2 + b^2$       (b)  $a^2 - b^2$   
 (c)  $a - b$       (d)  $a + b$
9. The degree of the polynomial  $x^2y^2 + 3xy + y^3$  is \_\_\_\_\_  
 (a) 4      (b) 5  
 (c) 6      (d) 2
10.  $x^2 - 4 =$  \_\_\_\_\_  
 (a)  $(x-2)(x+2)$       (b)  $(x-2)(x-2)$   
 (c)  $(x+2)(x+2)$       (d) None
11.  $x^3 + \frac{1}{x^3} = \left(x + \frac{1}{x}\right)(.....)$   
 (a)  $x^2 - 1 + \frac{1}{x^2}$       (b)  $x^2 + 1 + \frac{1}{x^2}$   
 (c)  $x^2 + 1 - \frac{1}{x^2}$       (d)  $x^2 - 1 - \frac{1}{x^2}$
12.  $2(a^2 + b^2) =$  \_\_\_\_\_  
 (a)  $(a+b)^2 + (a-b)^2$   
 (b)  $(a+b)^2 - (a-b)^2$   
 (c)  $(a+b)^2$       (d)  $4ab$
13. Order of surd  $\sqrt[3]{x}$  is \_\_\_\_\_  
 (a) 3      (b)  $\frac{1}{3}$   
 (c) 0      (d) 1

14.  $\frac{1}{2-\sqrt{3}} = \underline{\hspace{2cm}}$

- (a)  $2+\sqrt{3}$  (b)  $2-\sqrt{3}$   
(d)  $-2+\sqrt{3}$  (d)  $-2-\sqrt{3}$

15.  $(a+b)^2 - (a-b)^2 = \underline{\hspace{2cm}}$

- (a)  $2(a^2 + b^2)$  (b)  $4ab$   
(c)  $2ab$  (d)  $3ab$

16.  $\sqrt{14} \cdot \sqrt{35} = \underline{\hspace{2cm}}$

- (a)  $\sqrt[4]{10}$  (b)  $\sqrt[5]{10}$   
(c)  $7\sqrt{10}$  (d)  $8\sqrt{10}$

17. A surd which contains a single term is called surd.

- (a) Monomial  
(b) Binomial  
(c) Trinomial  
(d) None

### ANSWER KEY

1.	a	2.	d	3.	b	4.	a	5.	b
6.	b	7.	d	8.	c	9.	a	10.	a
11.	a	12.	a	13.	a	14.	a	15.	b
16.	c	17.	a						